Contribution ID: 29 Type: Poster

## Data analysis of the 26Si(a, p)29P reaction for the nucleosynthesis in the X-ray bursts

Monday, 9 September 2024 16:35 (1 minute)

The  $\alpha p$ -process, consisting of a series of  $(\alpha,p)$  and  $(p,\gamma)$  reactions, plays an important role in nucleosynthesis during X-ray bursts. Among these reactions, the  $26Si(\alpha,p)29P$  reaction is particularly significant due to its substantial impact on the light curve of X-ray bursts. Despite its importance, experimental data for this reaction remain insufficient due to technical reasons.

In this study, we performed a direct measurement of the  $26Si(\alpha,p)29P$  reaction at the CNS RI beam separator (CRIB). We measured the reaction at energies corresponding to the high-temperature environment around T = 3 GK using the 4He thick gas target method.

Despite numerous background events, we determined the reaction cross section, which was approximately an order of magnitude lower than the value of the NON-SMOKER statistical model.

This research represents the first experimental determination of this reaction cross section through direct measurement. Consequently, our results will provide valuable insights for estimating the reaction rate in the high-energy region and for refining the X-ray burst light curve model.

**Primary authors:** OKAWA, Kodai (CNS, the University of Tokyo); CHAE, Kyungyuk (Department of Physics, Sungkyunkwan University); KIM, Minju (Department of Physics, Sungkyunkwan University); HAYAKAWA, Seiya (Center for Nuclear Study, the University of Tokyo)

Co-authors: KIM, Chanhee (Department of Physics, Sungkyunkwan University); KIM, Dahee (Center for Exotic Nuclear Studies, Institute for Basic Science (IBS)); KAHL, David (Extreme Light Infrastructure Nuclear Physics (ELI-NP)); NGUYEN NGOC, Duy (Department of Physics, Sungkyunkwan University); GU, Gyungmo (Department of Physics, Sungkyunkwan University); SHIMIZU, Hideki (Center for Nuclear Study, the University of Tokyo); YA-MAGUCHI, Hidetoshi (Center for Nuclear Study, the University of Tokyo); LI, Jiatai (Center for Nuclear Study, the University of Tokyo); NGUYEN, KimUyen (Department of Physics, Sungkyunkwan University); SAKANASHI, Kohsuke (Department of Physics, Osaka University); KWAG, Minsik (Department of Physics, Sungkyunkwan University); MA, Nanru (Center for Nuclear Study, the University of Tokyo); IMAI, Nobuaki (Center for Nuclear Study, the University of Tokyo); SIRBU, Oana (Extreme Light Infrastructure Nuclear Physics (ELI-NP)); ZHANG, Qian (Center for Nuclear Study, the University of Tokyo); YOKOYAMA, Rin (Center for Nuclear Study, the University of Tokyo); ADACHI, Satoshi (Department of Physics, Osaka University); KUBONO, Shigeru (RIKEN Nishina Center); MICHIMASA, Shin'ichiro (Center for Nuclear Study, the University of Tokyo); HANAI, Shutaro (Center for Nuclear Study, the University of Tokyo); KIM, Sohyun (Department of Physics, Sungkyunkwan University); CHA, Soomi (Center for Exotic Nuclear Studies, Institute for Basic Science (IBS)); KAWABATA, Takahiro (Department of Physics, Osaka University); FURUNO, Tatsuya (Department of Physics, Osaka University); CHILLERY, Thomas (Center for Exotic Nuclear Studies, Institute for Basic Science (IBS))

**Presenter:** OKAWA, Kodai (CNS, the University of Tokyo)

Session Classification: Poster presentation

**Track Classification:** Experimental Nuclear Physics for Astrophysics