

# Structure Functions from the BK Equation Improved with Kinematical Constraint and DGLAP Evolution

Wanchen Li

wanchen1995@outlook.com

Department of Physics, Fudan University

In collaboration with:

Krzysztof Kutak, Anna Stasto, Robert Straka




**Light Cone 2024, Huizhou** — *November 24, 2024*

# Outline


- 1 Color Glass Condensate (CGC) and the Balitsky-Kovchegov (BK) Equation
  - Gluon saturation in the small  $x$  region
  - BK equation in the coordinate and momentum space
- 2 Resummations to the BK equation
  - Kinematical Constraints
  - The Dokshitzer–Gribov–Lipatov–Altarelli–Parisi (DGLAP) Splitting Functions
- 3 Structure Function from the  $k_T$  Factorization
- 4 Fit to Experimental Data
- 5 Conclusion

# Gluon saturation in the small $x$ region

- Gluon saturation: one of the major goals of the EIC to address.
- Ensuring unitarity for QCD or any non-Abelian theory.
- Initial condition for heavy-ion collisions, etc.



H1 and ZEUS, 2010.



Ciesielski, Kovchegov, Scapparone, DIS2015.


# Gluon saturation in the small $x$ region

- The strong rise (orange dashed line) in small  $k_T^2$  region, predicted by the Balitsky-Fadin-Kuraev-Lipatov (BFKL) equation violates the unitarity.
- When the transverse momentum  $k_T$  smaller than the saturation scale  $Q_s$ , that is

$$k^2 \equiv k_T^2 < Q_s^2(x).$$

The gluons start to recombine, regulating the infra-red divergence in the linear BFKL .

- The existence of energy dependent saturation scale is prediction of CGC.




Ciafaloni, Colferai, Salam, Stasto (CCSS)

WL, Stasto '22

## BK Equation in the Dipole Model

- The BK equation in the coordinate space:

$$\frac{\partial N(\mathbf{r}, \mathbf{b}, x)}{\partial \ln(1/x)} = \bar{\alpha}_s \int \frac{d^2 \mathbf{r}' r'^2}{(\mathbf{r}' + \mathbf{r})^2 (\mathbf{r}')^2} \left[ N\left(\mathbf{r}', \mathbf{b} + \frac{\mathbf{r}' + \mathbf{r}}{2}, x\right) + N\left(\mathbf{r}' + \mathbf{r}, \mathbf{b} + \frac{\mathbf{r}'}{2}, x\right) - N(\mathbf{r}, \mathbf{b}, x) - N\left(\mathbf{r}', \mathbf{b} + \frac{\mathbf{r}' + \mathbf{r}}{2}, x\right) N\left(\mathbf{r}' + \mathbf{r}, \mathbf{b} + \frac{\mathbf{r}'}{2}, x\right) \right]$$



Dipole scattering in DIS.

- $N(\mathbf{r}, \mathbf{b}, x)$  is the dipole amplitude,
- $\mathbf{r}$  is the size of the dipole,
- $\mathbf{b}$  is the impact parameter.
- The linear terms in the first two lines are BFKL terms
- The  $(N \cdot N)$  term in the last line corresponds to the non-linear evolution.

# BK Equation in the Momentum Space

- The BK equation in the momentum space:

Kutak, Kwiecinski '03; Nikolaev, Schafer '06; Bartels, Kutak '08.

$$\begin{aligned} \mathcal{F}(x, k^2) = & \mathcal{F}^{(0)}(x, k^2) + \bar{\alpha}_s \int_x^1 \frac{dz}{z} \int dk'^2 \left[ \frac{\mathcal{F}\left(\frac{x}{z}, k'^2\right)}{|k^2 - k'^2|} - \frac{k'^2}{k^2} \frac{2 \min(k^2, k'^2) \mathcal{F}\left(\frac{x}{z}, k\right)}{(k^2 + k'^2)|k^2 - k'^2|} \right] \\ & - \frac{2\bar{\alpha}_s^2 \pi^3}{N_c^2 R^2} \int_x^1 \frac{dz}{z} \left\{ \left[ \int_{k^2}^{\infty} \frac{dl^2}{l^2} \mathcal{F}\left(\frac{x}{z}, l^2\right) \right]^2 \right. \\ & \left. + \mathcal{F}\left(\frac{x}{z}, k^2\right) \int_{k^2}^{\infty} \frac{dl^2}{l^2} \ln\left(\frac{l^2}{k^2}\right) \mathcal{F}\left(\frac{x}{z}, l^2\right) \right\}. \end{aligned}$$

- The  $\mathcal{F}(x, k^2)$  is the unintegrated gluon density, related to the usual gluon PDF  $g(x, k^2)$  in double leading log approximation (DLLA) by

$$\mathcal{F}(x, k^2) = \frac{d}{dk^2} x g(x, k^2).$$

- The arguments of the running coupling  $\bar{\alpha}_s$  are associated with unintegrated gluon density respectively.
- $R$  is the size of the hadron's radius,  $N_c$  is the number of colors.

# Resummations

- To summarize the structure of our resummed equation, we give


$$\mathcal{F}(x, k^2) = \mathcal{F}^{(0)}(x, k^2) + \mathcal{K}_{\text{res}} \otimes \mathcal{F}(x, k^2) - \mathcal{V} \otimes \mathcal{F}^2(x, k^2).$$

- $\mathcal{F}^{(0)}(x, k^2)$  is the initial condition,
  - $\mathcal{V} \otimes \mathcal{F}^2(x, k^2)$  is the nonlinear term.
- $\mathcal{K}_{\text{res}} \otimes \mathcal{F}(x, k^2)$  is the resummed linear term

$$\mathcal{K}_{\text{res}} \otimes \mathcal{F} = \mathcal{K}_0^{\text{kc}}(z; \mathbf{k}, \mathbf{k}') \otimes^{z, \mathbf{q}} \mathcal{F}\left(\frac{x}{z}, k'\right) + \mathcal{K}_0^{\text{coll}}(z; k, k') \otimes^{z, k'} \mathcal{F}\left(\frac{x}{z}, k'\right),$$

- 'kc' stands for kinematical constraints,
- 'coll' stands for the collinear DGLAP evolution.

# Kinematical Constraints



Kwiecinski, Martin, Sutton '96

Andersson, Gustafson, Samuelsson '96

- The kinematical constraints can take different form due to the what approximation implemented, we use the first one below:

$$k'^2 < \frac{k^2}{z},$$

$$q^2 < \frac{k^2}{z},$$

$$q^2 < \frac{1-z}{z} k^2.$$

- The Kinematical constraints are due to that dominance of the transverse momentum in the small  $x$  region.





# Kinematical Constraints in the BFKL equation

- The kinematical constraint implement in the BFKL equation as

$$\begin{aligned} \mathcal{K}_0^{\text{kc}}(z; \mathbf{k}, \mathbf{k}') &\otimes^z \mathcal{F}\left(\frac{x}{z}, k'\right) \\ &= \int_x^1 \frac{dz}{z} \int \frac{d^2 \mathbf{q}}{\pi \mathbf{q}^2} \bar{\alpha}_s(\mathbf{q}^2) \left[ \mathcal{F}\left(\frac{x}{z}, |\mathbf{k} + \mathbf{q}|\right) \Theta\left(\frac{k^2}{z} - k'^2\right) - \Theta(k - q) \mathcal{F}\left(\frac{x}{z}, k\right) \right]. \end{aligned}$$

- $\chi_{\text{eff}}$  is the Pomeron intercept from the BFKL kernel in the Mellin space, indicating the power growth of the Reggeized cross section  $\sigma \sim s^{\chi_{\text{eff}}}$  with  $\chi_{\text{eff}}^{\text{LO BFKL}} = 4 \ln 2 \bar{\alpha}_s$ .



# Implementing DGLAP Splitting functions

- The resummation techniques that resums collinear and small  $x$  logarithms has been developed by several groups: Altarelli-Ball-Forte (ABF); Ciafaloni-Colferai-Salam-Stasto (CCSS), Thorne-White (TW).
- The DGLAP evolution is implemented as

$$\mathcal{K}_0^{\text{coll}}(z; k, k') \otimes_{z, k'} \mathcal{F}\left(\frac{x}{z}, k'\right) = \int_x^1 \frac{dz}{z} \int_0^{k^2} \frac{dk'^2}{k^2} \bar{\alpha}_s(k^2) z \tilde{P}_{gg}(z) \mathcal{F}\left(\frac{x}{z}, k'\right) \\ + \int_x^1 \frac{dz}{z} \int_{k^2}^{k'^2/z} \frac{dk'^2}{k'^2} \bar{\alpha}_s(k'^2) z \frac{k'^2}{k^2} \tilde{P}_{gg}\left(z \frac{k'^2}{k^2}\right) \mathcal{F}\left(\frac{x}{z}, k'\right).$$

- The two terms corresponds to the collinear and anti-collinear contribution, where the non-singular part of the splitting function is

$$\tilde{P}_{gg}^{(0)} = P_{gg}^{(0)} - \frac{1}{z},$$

# Structure Function from the $k_T$ Factorization


- The structure function  $F_2$  from the  $k_T$  factorization is given by

$$F_2(x, Q^2) = \sum_q e_q^2 S_q(x, Q^2),$$

where the sum is over the quark flavors and general expression for  $S_q(x, Q^2)$  is

$$S_q(x, Q^2) = \int_x^1 \frac{dz}{z} \int dk^2 S_{\text{box}}^q(z, m_q^2, k^2, Q^2) \mathcal{F}\left(\frac{x}{z}, k^2\right).$$

- $k$  is the gluon transverse momentum,  $\kappa$  is the quark transverse momentum.



DIS in the  $k_T$  factorization.

# Perturbative Contributions

- The off-shell photon-gluon partonic calculation gives

$$S_q(x, Q^2) = \frac{Q^2}{4\pi^2} \int \frac{dk^2}{k^2} \int_0^1 d\beta \int d\kappa' \alpha_s \left\{ [\beta^2 + (1 - \beta^2)] \left( \frac{\boldsymbol{\kappa}}{D_{1q}} - \frac{\boldsymbol{\kappa} - \mathbf{k}}{D_{2q}} \right)^2 + [m_q^2 + 4Q^2\beta^2(1 - \beta)^2] \left( \frac{1}{D_{1q}} - \frac{1}{D_{2q}} \right)^2 \right\} \mathcal{F}\left(\frac{x}{z}, k^2\right) \Theta\left(1 - \frac{x}{z}\right).$$

- The shifted quark transverse momentum is  $\boldsymbol{\kappa}' = \boldsymbol{\kappa} - (1 - \beta)\mathbf{k}$ .
- The energy denominators are

$$\begin{aligned} D_{1q} &= \kappa^2 + \beta(1 - \beta)Q^2 + m_q^2, \\ D_{2q} &= (\boldsymbol{\kappa} - \mathbf{k})^2 + \beta(1 - \beta)Q^2 + m_q^2. \end{aligned}$$

# Non-perturbative Contributions

- Depending on the values of transverse momenta  $k$ ,  $\kappa$  and a typical perturbative cut  $k_0 \sim 1$  GeV, following (Kwiecinski, Martin, Stasto '97), the perturbative and non-perturbative contributions can be categorized:

$$S_q = S_q^{(a)} + S_q^{(b)} + S_q^{(c)}.$$

- $S_q^{(a)}$ :  $k^2 < k_0^2$ ,  $\kappa^2 < k_0^2$ ; Modeled soft Pomeron contribution.
- $S_q^{(b)}$ :  $k^2 < k_0^2 < \kappa^2$ ; Previously modeled with collinear approximation.
  - We extend the lower bound  $k_{\min}^2 \ll k_0^2$  in the BK evolution to capture this contribution into the perturbative calculation, where the running coupling is frozen when  $k^2 < k_0^2$ .
- $S_q^{(c)}$ :  $k^2 > k_0^2$ ; Perturbative contribution from the  $k_T$  factorization.

# The Fit to the Structure Function

- We employ Golec-Biernat-Wuesthoff (GBW) inspired initial condition

$$\mathcal{F}^{(0)}(x, k) = A(1-x)^\alpha x^\beta \frac{\alpha_s(k^2)}{2} \left( k^{2\gamma_1} e^{-B_1^2 k^2} + \frac{1}{k^{2\gamma_2}} e^{-B_2^2/k^2} \right).$$


- The soft Pomeron contribution is parametrized as

$$S_q^{(a)} = C_{IP} x^{-\lambda} (1-x)^8.$$

- We fit to the HERA  $F_2$  data (2010), with  $\chi^2/\text{dof} = 1.54$ . Together with  $\Lambda_{\text{QCD}}$  and proton's radius  $R$ , the parameters are given as follows

Parameters	$\alpha$	$\beta$	$A$	$B_1$	$B_2$
Values	1.35	-0.425	1.245	0.436	1.50
$\gamma_1$	$\gamma_2$	$R^2$	$\Lambda_{\text{QCD}}$	$C_p$	$\lambda$
0.400	1.04	7.975	0.291	0.453	-0.0473


# The Fit to the Structure Function




- Fit describes a wide range of  $Q^2$ .

# The Fit to the Structure Function

BK without DGLAP



BK + DGLAP (our fit)




Dipole BK fits from Sanhueza, Carrido, Guevara, 2024.

- Including DGLAP on top of small  $x$  evolution enables a better fit for a wider range of  $Q^2$ .





# The Fit to the Reduced Cross Section



Fit to the HERA reduced cross sections (2015) yields  $\chi^2/\text{dof} = 1.82$ .

# The Unintegrated Gluon Density

Kutak, Sapeta '12; WL, Stasto '22.



- Our approach with an extensive range of  $k^2$  allows the extracting of the saturation scale  $Q_S^2$ .
- The saturation scale corresponds the  $k^2$  position for max unintegrated gluon density.
- One can see the  $Q_S^2$  increases when  $x$  decreases, as expected from the map of high energy QCD.

# Conclusion

- We implement the kinematical constraint and DGLAP resummation into the BK evolution.
- We achieve good fits to the structure functions and reduced cross sections, while the unintegrated gluon density is extended to low values of transverse momenta.
- Information of saturation scales can be extracted from our fit.
- Future Plans
  - Further implement the resummations in the dipole formalism.
  - Incorporating of the CCSS resummation with NLL BFKL into the framework.
  - Extend the study to the nuclei.