



Contribution ID: 57

Type: Oral

## Mass spectra of strange double charm pentaquarks with strangeness $S=-1$

Tuesday, 26 November 2024 16:50 (20 minutes)

The observation of the  $T_{c\bar{s}}(2900)$  indicates the potential existence of strange double charm pentaquarks based on the heavy antiquark symmetry. We systematically study the mass spectra of strange double charm pentaquarks with strangeness  $S = -1$  in both molecular and compact structures for quantum numbers  $J^P = 1/2^-, 3/2^-, 5/2^-$ . By constructing the interpolating currents, the mass spectra can be extracted from the two-point correlation functions in the framework of QCD sum rule method. In the molecular picture, we find that the  $\Xi_c'^+ D^{*+}$ ,  $\Xi_c^{*+} D^{*+}$ ,  $\Xi_{cc}^{*++} \bar{K}^{*0}$  and  $\Omega_{cc}^{*+} \rho^+$  may form molecular strange double charm pentaquarks. In both pictures, the masses of the  $J^P = 1/2^-, 3/2^-$  pentaquarks locate within the 4.2 – 4.6 GeV and 4.2 – 4.5 GeV regions, respectively. As all of them are above the thresholds of their strong decay channels, they behave as a broad state, making them challenging to be detected in experiment. On the contrary, the strange double charm pentaquark with  $J^P = 5/2^-$  lies below its strong decay channel, which may be a very narrow state and easy to be identified in experiment. The best observed channel is its semi-leptonic decay to double charm baryon. As the result, we strongly suggest experiments to search for  $J^P = 5/2^-$  strange double charm pentaquarks as a first try.

**Primary author:** Dr YANG, Zi-Yan

**Co-authors:** Prof. WANG, Qian (South China Normal University); Prof. CHEN, Wei (Sun Yat-sen University)

**Presenter:** Dr YANG, Zi-Yan

**Session Classification:** Parallel-3