Light-Cone 2024: Hadron Physics in the EIC era



Contribution ID: 62

Type: Oral

Application of Pionless Effective Field Theory in 3-hadron system.

Tuesday, 26 November 2024 17:30 (20 minutes)

Numerous low-lying excited hadrons can only decay into final states containing one or more photons, making them challenging to detect in experiments conducted at the Large Hadron Collider. Examples of such particles include the bottom partners of the $D_{s0}^*(2317)$ and $D_{s1}(2460)$, whose masses have been accurately predicted through heavy quark flavor symmetry and lattice calculations. We propose to measure the masses of such particles indirectly through quantum effects. Specifically, we aim to detect the threshold of a hadron and a companion hadron with attractive interactions using the threshold cusp effect. By subtracting the mass of the companion hadron from the threshold, we can determine the mass of the target hadron. We suggest detecting the thresholds of $B_{s0}^*\bar{B}_s$ and $B_{s1}\bar{B}_s$ in the energy distribution. Furthermore, we demonstrate that the masses of $D_{s0}^*(2317)$ and $D_{s1}(2460)$ can be extracted from the J/ distribution measured by the LHCb Collaboration.

Primary author: FU, Hailong (ITP)

Co-authors: Prof. HANHART, Christoph (FZ Julich, IKP); GUO, Feng-Kun (ITP, CAS); Dr YAN, Maojun (Southwest University); Dr ZHANG, Xu (ITP, CAS)

Presenter: FU, Hailong (ITP)

Session Classification: Parallel-3